

Energy conditions and a bounce in FLRW cosmologies

Benjamin K. Tippett and Kayll Lake [*]

Department of Physics, Queen's University, Kingston, Ontario, Canada, K7L 3N6

(Dated: February 3, 2008)

Exploiting the existence of two “cosmological” constants, one associated with the classical Lovelock theorem and one with the vacuum energy density, we argue, in a model independent way, that in spatially closed FLRW cosmologies with a positive definite effective cosmological constant there exists a range in this constant that serves as a sufficient condition for the satisfaction of the null, weak, strong and dominant energy conditions at a bounce. The application of energy conditions is not unambiguous and we show how the bounce can be considered classically and how, we believe more reasonably, it can be considered a matter of quantum cosmology.

Introduction. The idea of a bounce in Friedmann-Lemaître-Robertson-Walker (FLRW) cosmologies is an idea that is as old as spacetime cosmology itself [1] but also one with current appeal [2] motivated primarily by M-theory, branes and extra-dimensions [3]. A bounce, as opposed to a big bang, obviously meets with many problems when trying to reproduce the universe we observe. Moreover, there is a widely held belief that a bounce must violate classical energy conditions [4]. However, the application of classical energy conditions is not unambiguous and here we exploit the existence of two “cosmological” constants, one associated with the classical Lovelock theorem and one with the vacuum energy density, and argue, in a model independent way, that in spatially closed FLRW cosmologies with a positive definite effective cosmological constant (as explained precisely below) there exists a range in this constant that serves as a sufficient condition for the satisfaction of the null, weak, strong and dominant energy conditions at a bounce. Further, depending on how the energy conditions are applied, we show how a bounce might be considered classically and how, we believe more correctly, it must be considered a matter of quantum cosmology. In this analysis we offer no explanation as to why there should be a bounce but rather only examine the consequences of a bounce as regards the application of energy conditions.

The suggestion that we live in a spatially closed universe with a positive definite observed cosmological constant is in accord with the recent Hubble Space Telescope observations of Riess *et al.* [5], observations that not only confirm earlier reports that we live in an accelerating universe [6] [7], but also sample the transition from deceleration to acceleration. In contrast, observations with the Wilkinson Microwave Anisotropy Probe [8] argue strongly that the FLRW cosmology is spatially flat [9]. In the spatially flat case we argue that a bounce necessarily violates energy conditions no matter how they are applied. Because our application of the energy conditions differs in a significant way from previous works, we start our discussion from first principles.

Background Geometry. The Cosmological Principle is the working hypothesis that the Universe is described by a spacetime that is spatially isotropic globally. It follows from the theorems of Robertson and of Walker [10]

that the spacetime is represented by the Friedmann-Lemaître-Robertson-Walker (FLRW) metric, which, up to a coordinate transformation, is given by [11]

$$ds^2 = a^2 d\bar{s}_\Gamma^2 - dt^2 \quad (1)$$

where $a = a(t)$, normally $\in C^2$, and $d\bar{s}_\Gamma^2$ is the metric of a three space of constant curvature. In Γ we scale the coordinates so that $k \equiv R_a^b R_b^a / 2(R_a^a)^2 = \pm 1, 0$ where R_a^b is the Ricci tensor of Γ .

The metric (1) has only two independent (scalar) invariants derivable from the Riemann tensor without differentiation [12]. These are usually taken to be the Ricci scalar R ($\equiv R_\alpha^\alpha$, R_α^β the Ricci tensor of the spacetime) and first Ricci invariant $R1$ ($\equiv S_\alpha^\beta S_\beta^\alpha / 4 \equiv S^2$, S_α^β the trace-free Ricci tensor $\equiv R_\alpha^\beta - \delta_\alpha^\beta R/4$). Without any loss in generality we can also take the two independent invariants to be ρ and p defined as follows:

$$8\pi\rho \equiv R/4 + \text{sign}(S)\sqrt{3S^2} - \mathcal{C} \quad (2)$$

and

$$8\pi p \equiv -R/4 + \text{sign}(S)\sqrt{S^2/3} + \mathcal{C}, \quad (3)$$

where $\mathcal{C}$ is a constant. In the coordinates of (1) these evaluate to

$$8\pi\rho = \frac{3}{a^2}(k + \dot{a}^2) - \mathcal{C}, \quad (4)$$

and

$$8\pi p = -\frac{1}{a^2}(k + \dot{a}^2 + 2a\ddot{a}) + \mathcal{C}, \quad (5)$$

respectively where $\cdot \equiv d/dt$. Since ρ and p are invariants, they distinguish scalar polynomial singularities of the metric (1) directly [13].

Physical Interpretation. A remarkable feature of four dimensions is the fact that the only divergence free two index tensor A_β^α derivable from the metric tensor and its first two derivatives is (up to a disposable multiplicative constant) [14]

$$A_\beta^\alpha = G_\alpha^\beta + \lambda_c \delta_\alpha^\beta, \quad (6)$$

where G_α^β is the Einstein tensor ($\equiv R_\alpha^\beta - \frac{1}{2}R\delta_\alpha^\beta$) and λ_c is a constant. We write Einstein's equations in the form

$$G_\alpha^\beta + \lambda_c \delta_\alpha^\beta = 8\pi_M T_\alpha^\beta + 8\pi_V T_\alpha^\beta, \quad (7)$$

where $_M T_\alpha^\beta$ signifies the matter contribution to (1) and $_V T_\alpha^\beta$ the vacuum contribution. Here we simply assume that the latter is of the usual form $8\pi_V T_\alpha^\beta \equiv \lambda_v \delta_\alpha^\beta$ where λ_v a constant. Central to our considerations here is the treatment of the constant λ_c which, it is fair to say, is usually ignored.

It is useful at this point, prior to the application of energy conditions, to review three distinct ways of proceeding with (7) [15]. Because of the form of (1) we can always consider the source a perfect mathematical fluid (including, for example, scalar fields [16]) with associated streamlines comoving with Γ [17]. The flow (congruence of unit timelike vectors u^α) and normal field n^α (in the tangent space of the associated Lorentzian two-space) are uniquely determined. The energy density and isotropic pressure (including bulk viscosity) are now defined by

$$\rho \equiv T_\alpha^\beta u^\alpha u_\beta \quad (8)$$

and

$$p \equiv T_\alpha^\beta n^\alpha n_\beta \quad (9)$$

respectively where T_α^β is the energy-momentum tensor of the mathematical fluid. The definitions (8) and (9) reproduce (4) and (5) (now recognized as the Friedmann equations [18]) from the field equations

$$G_\alpha^\beta + \mathcal{C} \delta_\alpha^\beta = 8\pi T_\alpha^\beta. \quad (10)$$

The question of interest here is the treatment of the constants λ_c and λ_v in the procedure of applying the energy conditions. The three distinct ways of proceeding are:

(i) $\mathcal{C} = \lambda_c - \lambda_v \equiv \Lambda$: First note that Λ is an observable (let us call it the observed cosmological constant). The apparent smallness of Λ arises out of the almost perfect cancellation of λ_c and λ_v an explanation of which constitutes a standard form of one of the ‘‘cosmological constant problems’’ [19]. The procedure of setting $\mathcal{C} = \lambda_c - \lambda_v \equiv \Lambda$ is the common one when energy conditions are not under consideration and, for example, is the procedure used (implicitly) to construct the observers $\Omega_\Lambda - \Omega_M$ plane [9]. As regards the application of energy conditions, this procedure is *ad hoc* since the vacuum contribution is, without explanation, extracted from the energy-momentum tensor and considered part of the geometry. We write the associated energy density and isotropic pressure of the perfect fluid source as ρ and p respectively. These are given by (4) and (5). (We view $\mathcal{C} = -\lambda_v$ as a special case where one simply assumes, without explanation, that $\lambda_c = 0$.)

(ii) $\mathcal{C} = \lambda_c$: In this procedure the vacuum contribution is not extracted from the energy-momentum tensor but

rather considered part of it. The constant λ_c is not directly observable. Rather, its value can only be inferred from the fact that $\lambda_c \sim \lambda_v$ and with λ_v calculated as in the standard model. The perfect fluid source now has an associated energy density and isotropic pressure given by $\tilde{\rho} \equiv \rho - \frac{\lambda_v}{8\pi}$ and $\tilde{p} \equiv p + \frac{\lambda_v}{8\pi}$ respectively. These are given by (4) and (5) with ρ and p replaced by $\tilde{\rho}$ and $\tilde{p}$. This procedure, as regards the application of energy conditions, is clearly more natural than (i).

(iii) $\mathcal{C} = 0$: In this procedure the geometrical contribution from λ_c and the contribution from the vacuum λ_v are both considered part of the energy-momentum tensor. Exactly why the geometrical contribution λ_c should be considered part of energy-momentum tensor is left unexplained. The perfect fluid source now has an associated energy density and isotropic pressure given by $\bar{\rho} \equiv \rho - \frac{\lambda_v - \lambda_c}{8\pi}$ and $\bar{p} \equiv p + \frac{\lambda_v - \lambda_c}{8\pi}$ respectively. Again these are given by (4) and (5) with ρ and p replaced by $\bar{\rho}$ and $\bar{p}$. (This procedure can also be considered as a special case of (ii) where one simply assumes, again without explanation, that $\lambda_c = 0$.)

Energy Conditions. We now impose standard energy conditions on the mathematical fluid [20] [21] [22]. The local energy conditions considered here are the null energy condition (NEC), weak energy condition (WEC), strong energy condition (SEC) and dominant energy condition (DEC) [23]. We let EC designate the simultaneous satisfaction of all four energy conditions. In the present context we are always dealing with a perfect fluid and in that case the conditions are given in case (i) by

$$\text{NEC} \iff (\rho + p \geq 0), \quad (11)$$

$$\text{WEC} \iff (\rho \geq 0) \text{ and } (\rho + p \geq 0), \quad (12)$$

$$\text{SEC} \iff (\rho + 3p \geq 0) \text{ and } (\rho + p \geq 0), \quad (13)$$

and

$$\text{DEC} \iff (\rho \geq 0) \text{ and } (\rho \pm p \geq 0), \quad (14)$$

where ρ and p are replaced by $\tilde{\rho}$ and $\tilde{p}$ in case (ii) and by $\bar{\rho}$ and $\bar{p}$ in case (iii). From (4) and (5) for case (i) we find

$$\rho + 3p \geq 0 \iff \mathcal{C} \geq \frac{3\ddot{a}}{a}, \quad (15)$$

$$\rho + p \geq 0 \iff k + \dot{a}^2 \geq a\ddot{a}, \quad (16)$$

$$\rho - p \geq 0 \iff \mathcal{C} \leq \frac{2}{a^2}(k + \dot{a}^2) + \frac{\ddot{a}}{a}, \quad (17)$$

and

$$\rho \geq 0 \iff \mathcal{C} \leq \frac{3}{a^2}(k + \dot{a}^2), \quad (18)$$

again with ρ and p are replaced by $\tilde{\rho}$ and $\tilde{p}$ in case (ii) and by $\bar{\rho}$ and $\bar{p}$ in case (iii). Case (iii) has been considered in detail previously [4]. As we show below, this case is not of interest for our present considerations. From the foregoing conditions it follows that [24]

$$\frac{3\ddot{a}}{a} \leq \mathcal{C} \leq \frac{2}{a^2}(k + \dot{a}^2) + \frac{\ddot{a}}{a} \Rightarrow \text{EC.} \quad (19)$$

A bounce. We characterize regular minima via the conditions

$$\dot{a} = 0, \quad \ddot{a} > 0, \quad a > 0 \quad (20)$$

at, say, $t = t_m$. It follows immediately from (16) that $k = 1$ at a regular minimum (and in particular, $k = 0$ is not a possibility). Whereas the timelike convergence condition ($R^\alpha_\beta W^\alpha W^\beta \geq 0 \quad \forall$ timelike W^α) fails at all regular minima (e.g. $W^\alpha = u^\alpha$), it follows from (19) that there remains in general a non-vanishing interval in $\mathcal{C}$ sufficient for the satisfaction of the energy conditions in cases (i) with $\lambda_c > \lambda_v$ and (ii) with $\lambda_c > 0$. Indeed with $k = 1$ and (20) it follows that the range in $\mathcal{C}$ can be narrowed to give

$$\frac{3\ddot{a}}{a} \leq \mathcal{C} \leq \frac{2}{a^2} \Rightarrow \text{EC.} \quad (21)$$

The marginal case $\dot{a} = \ddot{a} = 0, \quad a > 0$ distinguishes flatness at $t = t_m$ ($k = \mathcal{C} = 0$) and $k = 1$ with the obvious substitutions in (19) and (21). In both these cases the timelike convergence condition holds [25].

Discussion. We have shown, in a model independent way, that in a spatially closed FLRW universe there exists a range in $\mathcal{C} > 0$ in cases (i) and (ii) that serves as a sufficient condition for the satisfaction of the standard energy conditions at a bounce. Since we have used no properties of the scale factor a , other than the existence of a bounce, nothing can be said about the lower limit

$$\mathcal{C} \geq \frac{3\ddot{a}}{a}|_{t_m} > 0. \quad (22)$$

Expressing the upper limits in terms of $a(t_m)$ we have

$$a(t_m) \leq \sqrt{\frac{2}{\Lambda}}, \quad a(t_m) \leq \sqrt{\frac{2}{\lambda_c}} \quad (23)$$

in cases (i) and (ii) respectively. In case (i), since $\Lambda < \sim 10^{-56} \text{cm}^{-2}$ the lower bound on the upper limit to $a(t_m)$ is of the order of $4.5 \cdot 10^3$ Mpc essentially equivalent to the current Hubble scale ($\frac{c}{H_0}$) and so one might hope that the existence or not of a bounce might be answerable without recourse to quantum cosmology. The fact that the energy conditions need not be violated at a bounce for cosmologies consistent with the Hubble Space Telescope observations is certainly of interest. However, as pointed out above, this procedure, as regards the application of energy conditions, is *ad hoc* in that it extracts λ_v from the energy-momentum tensor without any explanation.

In case (ii) we must *assume* that $\lambda_c > 0$ for the energy conditions not to be violated at a bounce. Moreover, in contrast to case (i), if we trust the standard model all the way to the Planck scale, then the lower bound on the upper limit to $a(t_m)$ is within an order or so of the Planck scale itself and the existence or not of a bounce becomes a matter of quantum cosmology. This procedure has to be considered the preferred one since the role of λ_c and of λ_v is unaltered by *ad hoc* assumptions. Whereas the classical energy conditions may not play a fundamental role in quantum cosmology, the fact that they need not be violated at a bounce is certainly of interest. Moreover, the avoidance of energy condition violations at a bounce is tied to the mysterious cosmological constant problem wherein the purely classical statement (6) is, by the observation that $\lambda_c \sim \lambda_v$, intimately connected to the non-classical calculation of λ_v .

Acknowledgments

This work was supported by a grant to KL from the Natural Sciences and Engineering Research Council of Canada.

[*] Electronic Address: lake@astro.queensu.ca

- [1] Two classic examples are the discussions in the text by Tolman (R.C. Tolman, *Relativity, Thermodynamics and Cosmology*, (Oxford Univ. Press, London, 1934)) and in the review by Robertson (H. P. Robertson, *Rev. Mod. Phys.*, **5**, 62 (1933)).
- [2] For example, a search at arXiv.org of “bouncing cosmologies” returns over 100 abstracts at the time of writing.
- [3] See, for example, P. J. Steinhardt and N. Turok, “The Cyclic Universe: An Informal Introduction” (astro-ph/0204479).
- [4] See C. Molina-Paris and M. Visser, *Phys. Lett. B*, **455**, 90 (1999) (gr-qc/9810023) and, for a less general analysis,

C.P. Burgess, F. Quevedo, R. Rabadan, G. Tasinato and I. Zavala, *JCAP* 0402 (2004) 008. (hep-th/0310122)

- [5] A. G. Riess *et al.*, “Type Ia Supernova Discoveries at $z > 1$ From the Hubble Space Telescope: Evidence for Past Deceleration and Constraints on Dark Energy Evolution”, *Ap. J.* **607**, 665 (2004). (astro-ph/0402512)
- [6] A. G. Riess *et al.*, *A. J.*, **116** 1009 (1998). (astro-ph/9805201)
- [7] S. Perlmutter *et al.*, *Ap. J.*, **517**, 565 (1999). (astro-ph/9812133)
- [8] D. N. Spergel *et al.*, “First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters”, *Ap. J. Suppl.* **148**,

- 175 (2003) (astro-ph/0302209)
- [9] See, for example, K. Lake, “Cosmological observations and their representation in the $\Omega_\Lambda - \Omega_M$ plane” (2004). (astro-ph/0404319)
- [10] The uniqueness of the FLRW metric can be derived without any reference to field equations and in particular without reference to any background mathematical fluid. In this regard it is perhaps useful to view the metric (1) as derived from exact spherical symmetry about every point. See A. G. Walker, J. London. Math. Soc., **19**, 219 (1944).
- [11] We use a signature of +2 and start with geometrical units. Latin indicies range from 1 to 3 and Greek indicies from 1 to 4.
- [12] See, for example, K. Santosuosso, D. Pollney, N. Pelavas, P. Musgrave and K. Lake, Computer Physics Communications **115**, 381 (1998). (gr-qc/9809012)
- [13] In particular, there is no need to look at the traditional “Kretschmann” invariant $K = R_{\alpha\beta\gamma\delta}R^{\alpha\beta\gamma\delta}$ ($R_{\alpha\beta\gamma\delta}$ the Riemann tensor).
- [14] D. Lovelock, J. Math. Phys. **13**, 874 (1972).
- [15] For use of these distinctions in a different context see K. Lake, “Sudden future singularities in FLRW cosmologies”, Classical and Quantum Gravity (in press) (gr-qc/0407107)
- [16] See, for example, *Dynamical Systems in Cosmology* Edited by J. Wainwright and G. F. R. Ellis (Cambridge University Press, Cambridge, 1997).
- [17] For further discussion see H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Herlt, *Exact Solutions of Einstein's Field Equations* (Cambridge University Press, Cambridge, 2003) and A. Krasinski, *Inhomogeneous Cosmological Models* (Cambridge University Press, Cambridge, 1997).
- [18] Friedmann's papers, with expert and referenced editorial notes, have been reprinted in English translation in *Gen. Rel. Grav.* **31**, 1985 (1999).
- [19] See, for example, S. Weinberg astro-ph/9610044 and citations thereto and T. Padmanabhan, Phys.Rept. **380**,235 (2003). (hep-th/0212290).
- [20] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, 1973)
- [21] M. Visser, *Lorentzian Wormholes* (Springer-Verlag, New York, 1996)
- [22] E. Poisson, *A Relativist's Toolkit: The Mathematics of Black-hole Mechanics* (Cambridge University Press, Cambridge, 2004).
- [23] These conditions are usually stated as inequities that hold for any vector field of a stated type. By this we understand that the inequalities must hold over the set of all possible vector fields of a stated type. The inequalities are in general not appropriate for a subset as the associated physics can be entirely regular for the subset under inequalities that differ from the stated conditions.
- [24] Note that the existence of this interval (without $\mathcal{C}$) is required if the spacetime is to satisfy the null energy condition.
- [25] At regular maxima consider $\ddot{a} < 0$ in the foregoing and it follows that only for $k = 1$ can $\mathcal{C}$ be > 0 in the sufficient condition for EC.